

Fractalkine (CX3CL1) and brain inflammation: Implications for HIV-1-associated dementia

R Cotter,^{1,2} C Williams,^{1,2} L Ryan,^{1,2} David Erichsen,^{1,2} A Lopez,^{1,2} H Peng,^{1,2} and J Zheng^{1,2,3}

¹The Laboratory of Neurotoxicology, the Center for Neurovirology and Neurodegenerative Disorders, and the Departments of ²Pathology and Microbiology and ³Pharmacology, University of Nebraska Medical Center, Omaha, Nebraska, USA

Leukocyte migration and activation play an important role in immune surveillance and the pathogenesis of a variety of neurodegenerative disorders, including human immunodeficiency virus (HIV)-1-associated dementia (HAD). A novel chemokine named fractalkine (FKN, CX3CL1), which exists in both membrane-anchored and soluble isoforms, has been proposed to participate in the generation and progression of inflammatory brain disorders. Upon binding to the CX3C receptor one (CX3CR1), FKN induces adhesion, chemoattraction, and activation of leukocytes, including brain macrophages and microglia (MP). Constitutively expressed in the central nervous system (CNS), mainly by neurons, FKN is up-regulated and released in response to proinflammatory stimuli. Importantly, FKN is up-regulated in the brain tissue and cerebrospinal fluid (CSF) of HAD patients. Together, these observations suggest that FKN and its receptor have a unique role in regulating the neuroinflammatory events underlying disease. This review will examine how FKN contributes to the recruitment and activation of CX3CR1-expressing MP, which are critical events in the neuropathogenesis of HAD. *Journal of NeuroVirology* (2002) **8**, 585–598.

Keywords: chemokine receptors; chemokines; fractalkine; HIV-1-associated dementia

Introduction

Human immunodeficiency virus (HIV)-1-associated dementia (HAD) is a late-stage complication of advanced HIV-1 disease (Carpenter *et al*, 2000; Krebs *et al*, 2000; McArthur *et al*, 1999). Clinically, HAD results in a spectrum of neurological and psychiatric symptoms, including cognitive impairment, hallucinations, delirium, coma, and ultimately death (Gelbard and Epstein, 1995; Janssen *et al*, 1991; Marder *et al*, 1996; Masliah, 1996; Navia *et al*, 1986).

The histopathological correlate of HAD is HIV-1 encephalitis (HIVE), which occurs in most, but not all, cases of dementia related to HIV-1 infection (Glass *et al*, 1995; Masliah, 1996; Wiley, 1995). HIVE features blood-brain-barrier (BBB) damage, productive viral infection, immune activation of mononuclear phagocytes (MP; brain macrophage and microglia), astrogliosis, and neuronal injury, apoptosis, and loss (Asare *et al*, 1996; Dickson *et al*, 1994; Gabuzda and Wang, 1999; Gendelman *et al*, 1997; Glass *et al*, 1995; Masliah *et al*, 2000; McArthur *et al*, 1999; Nath and Geiger, 1998; Navia *et al*, 1986; Rappaport *et al*, 1999; Wiley and Achim, 1994). It is believed that MP, the predominate cell type infected in the brain, induce neuronal injury and death through the production of neurotoxins (Gendelman *et al*, 1997; Genis *et al*, 1992; Giulian *et al*, 1990; Koenig *et al*, 1986; Ma *et al*, 1994; Moses *et al*, 1993; Nath *et al*, 1995; Pulliam *et al*, 1991; Ranki *et al*, 1995; Tornatore *et al*, 1991; Wiley *et al*, 1991). Given what is known about the involvement of MP in HIVE, it is important to understand how MP become immune activated. Recently, it has been proposed that neurons may directly participate in the disease process by inducing MP

Address correspondence to Jialin Zheng, Laboratory of Neurotoxicology, Center for Neurovirology and Neurodegenerative Disorders, 985215 Nebraska Medical Center, Omaha, NE 68198-5215, USA. E-mail: jzheng@unmc.edu

The authors kindly thank Dr. Howard E Gendelman, Mr. Douglas Niemann, and Mr. Michael Bauer for scientific discussion and suggestions; and Ms. Julie Ditter, Ms. Robin Taylor, and Ms. Theresa Grutel for outstanding administrative and secretarial support. This work was supported in part by research grants P20 RR15635-01 and R01 NS 41858-01 from the National Institutes of Health (to JZ).

Received 15 February 2002; revised 13 May 2002; accepted 1 July 2002.

recruitment and activation through release of soluble chemotactic factors. This review will examine the role that neuronal chemokines play in MP recruitment and activation during HAD.

Chemokines and chemokine receptors in the CNS

Chemoattractant cytokines (chemokines) are soluble molecules that regulate the migration and activation of leukocytes into brain and other tissues (Kutsch *et al*, 2000; Wu *et al*, 2000). More than 46 chemokines have been identified (Baggiolini *et al*, 1997; Zlotnik and Yoshie, 2000) and are classified into the following four groups (Table 1): alpha (CXC), beta (CC), gamma (C-chemokines), and delta (CX3C), based on the arrangement of cysteine residues within the receptor-binding domain. For example, CXC chemokines have two cysteine residues separated by a single amino acid, whereas in CC chemokines the cysteines are adjacent. Chemokines exert their effects by binding to and activating a family of seven-transmembrane, G-protein-coupled receptors (GPCRs). These receptors are divided into four groups: α -chemokine receptors (CXCR1-6), β -chemokine receptors (CCR1-10), γ -chemokine receptors (XCR1), and δ -chemokine receptors (CX3CR1) (Hesseltger and Horuk, 1999; Klein *et al*, 1999; van der Meer *et al*, 2000) (review in Gabuzda *et al*, 2002; Karpus, 2001; Miller and Meucci, 1999; Ransohoff, 1998). In addition to mediating leukocyte recruitment and activation, chemokine receptors, such as CCR5 and CXCR4, also serve as coreceptors for HIV-1 (Dragic *et al*, 1996; He *et al*, 1997). Importantly, the endogenous ligands (RANTES, macrophage inflammatory protein [MIP]-1 α/β) for these receptors have been shown to block HIV-1 binding and entry, suggesting that the production of these factors may be an important defense mechanism against HIV-1 infection in the human host (Kornbluth *et al*, 1998).

A wide range of chemokines are expressed in the brain during diseases, including α -chemokines, such as interleukin-8 (IL-8, CXCL8) and stromal-derived factor-1 alpha (SDF-1 α , CXCL12); β -chemokines, such as monocyte chemoattractant protein (MCP)-1 (CCL2), MIP-1 α (CCL3), MIP-1 β (CCL4), and RANTES (CCL5); and the δ -chemokine, fractalkine (FKN, CX3CL1) (Conant *et al*, 1998; Cotter *et al*, 1999b; Coughlan *et al*, 2000; Desbaillets *et al*, 1994; Gabuzda and Wang, 2000; Kornbluth *et al*, 1998; Persidsky, 1999; Zheng *et al*, 1999). Chemokines, such as SDF-1 α , IL-8, and fractalkine (FKN), are constitutively produced in the brain and play an important role in central nervous system (CNS) homeostasis and development (Coughlan *et al*, 2000; Gabuzda and Wang, 2000; Gleichmann *et al*, 2000; Harrison *et al*, 1998; Horuk *et al*, 1996; Meucci *et al*, 1998, 2000; Nagasawa *et al*, 1996). Upon binding to neu-

Table 1 Chemokine and chemokine receptor families*

| Systematic name | Human ligand | Chemokine receptors |
|--|------------------------------------|---------------------|
| Alpha (CXC) chemokine-receptor family | | |
| CXCL1 | GRO α /MGS α | CXCR2, CXCR1 |
| CXCL2 | GRO β /MGS β | CXCR2 |
| CXCL3 | GRO γ /MGS γ | CXCR2 |
| CXCL4 | PF4 | Unknown |
| CXCL5 | ENA-78 | CXCR2 |
| CXCL6 | GCP-2 | CXCR1, CXCR2 |
| CXCL7 | NAP-2 | CXCR2 |
| CXCL8 | IL-8 | CXCR1, CXCR2 |
| CXCL9 | Mig | CXCR3 |
| CXCL10 | IP-10 | CXCR3 |
| CXCL11 | I-TAC | CXCR3 |
| CXCL12 | SDF-1 (α/β) | CXCR4 |
| CXCL13 | BCA-1 | CXCR5 |
| CXCL14 | BRAK/bolkinge | Unknown |
| (CXCL15) | Unknown | Unknown |
| CXCL16 | Unknown | CXCR6 |
| Beta (CC) chemokine-receptor family | | |
| CCL1 | I-309 | CCR8 |
| CCL2 | MCP-1/MCAF/TDCF | CCR2 |
| CCL3 | MIP-1 α /LD78 α | CCR1, CCR5 |
| CCL4 | MIP-1 β | CCR5 |
| CCL5 | RANTES | CCR1, CCR3, CCR5 |
| (CCL6) | Unknown | Unknown |
| CCL7 | MCP-3 | CCR1, CCR2, CCR3 |
| CCL8 | MCP-2 | CCR3, CCR5 |
| (CCL9/10) | Unknown | CCR1 |
| CCL11 | Eotaxin | CCR3 |
| (CCL12) | Unknown | CCR2 |
| CCL13 | MCP-4 | CCR2, CCR3 |
| CCL14 | HCC-1 | CCR1, CCR5 |
| CCL15 | HCC-2/Lkn-1/MIP-1 δ | CCR1, CCR3 |
| CCL16 | HCC-4/LEC/LCC-1 | CCR1, CCR2 |
| CCL17 | TARC | CCR4 |
| CCL18 | DC-CK1/PARC/AMAC-1 | Unknown |
| CCL19 | MIP-3 β /ELC/exodus-3 | CCR7 |
| CCL20 | MIP-3 α /LARC/exodus-1 | CCR6 |
| CCL21 | 6Ckine/SLC/exodus-2 | CCR7 |
| CCL22 | MDC/STCP-1 | CCR4 |
| CCL23 | MPIF-1/CK β 8/CK β 8-1 | CCR1 |
| CCL24 | Eotaxin-2/MPIF-2 | CCR3 |
| CCL25 | TECK | CCR9 |
| CCL26 | Eotaxin-3 | CCR3 |
| CCL27 | CTACK/ILC | CCR10 |
| CCL28 | MEC | CCR3/CCR10 |
| Gamma (C) Chemokine/receptor family | | |
| XCL1 | Lymphotactin/SCM-1 α | XCR1 |
| XCL2 | SCM-1 β | XCR1 |
| Delta (CX3C) chemokine/receptor family | | |
| CX3CL1 | Fractalkine/neurotactin | CX3CR1 |

*Modified from Zlotnik and Yoshie (2000).

ronal receptors (CXCR4, CXCR2, and CX3CR1), these chemokines activate signaling pathways that regulate neuronal survival, injury, and repair (Kaul and Lipton, 1999; Meucci *et al*, 1998, 2000; Peng *et al*, 2002; Tong *et al*, 2000; Zheng *et al*, 2001). For example, knockout mice lacking CXCR4 exhibit abnormal migration of cerebellar external granule layer cells and other nervous system defects (Zou *et al*, 1998). These findings underscore the importance of chemokines and their receptors in neuronal cell development and maintenance.

Chemokines also play a critical role in the host response to CNS injury and infection. Indeed, the role of chemokines and their receptors in neurodegenerative disorders, such as multiple sclerosis, Alzheimer's disease, stroke, and HAD, has been extensively investigated and reviewed (Gabuzda *et al*, 1998, 2002; Karpus, 2001; Letendre *et al*, 1999; Minami and Satoh, 2000; Ransohoff, 1997; Sanders *et al*, 1998). Several reports have shown that SDF-1 α , IL-8, MIP-1 α , MIP-1 β , RANTES, MCP-1, and FKN are up-regulated in brain tissue and cerebrospinal fluid (CSF) from HAD patients (Coughlan *et al*, 2000; Kelder *et al*, 1998; Persidsky *et al*, 1999; Tong *et al*, 2000; Zheng *et al*, 1999, 2000). It has been proposed that these chemokines contribute to HAD pathogenesis through recruitment of monocytes into the brain, through initiation of neuroinflammatory cascades that affect viral replication, through induction of neural signaling and apoptosis, or through initiation of neuronal protection and repair (Albright *et al*, 1999; Broder and Collman, 1997; Cotter *et al*, 2001; Endres *et al*, 1996; Ghorpade *et al*, 1998; He *et al*, 1997; Kitai *et al*, 2000; Lavi *et al*, 1997; Luster, 1998; Mackay, 1996; Shieh *et al*, 1998; Vallat *et al*, 1998; Vicenzi *et al*, 2000; Zheng *et al*, 1999) (review in Gabuzda *et al*, 2002; Miller and Meucci, 1999). Although it is evident that chemokines are an important component of the host immune response, the nature of their role in disease pathogenesis is only beginning to be understood.

Neuronal chemokines and MP activation, a "chicken or egg" question

Traditionally, it was believed that MP activation and chemokine production preceded neuronal injury in HIVE. However, new evidence suggests that neurons themselves may initiate MP recruitment and activation (Biber *et al*, 2001; Harrison *et al*, 1998). Indeed, it has been proposed that in response to injury, neurons produce chemokines, such as FKN (Harrison *et al*, 1998), that act as "distress signals." Upon release, these factors recruit MP to sites of injury and stimulate the production of inflammatory factors with the potential to repair or exacerbate neuronal damage (Tong *et al*, 2000; Zheng *et al*, 2000) (Figure 1).

There are several lines of evidence that provide support for this hypothesis. *First*, excitotoxin-mediated neuronal injury, as well as nerve axotomy, induces production of the neuronal chemokine FKN (Chapman *et al*, 2000a; Harrison *et al*, 1998; Zheng *et al*, 2000). *Second*, this neuronal chemokine is also up-regulated in HAD brain tissue (Pereira *et al*, 2001; Tong *et al*, 2000; Zheng *et al*, 2000) and released in response to neuronal apoptosis induced by HIV-1 progeny virions (IIIB and ADA) and gp120 (Zheng *et al*, 2000). *Third*, both the soluble and membrane-bound forms of FKN have been shown to attract and immobilize leukocytes, such as monocytes and lym-

phocytes (Boehme *et al*, 2000; Chapman *et al*, 2000a, 2000b; Combadiere *et al*, 1998; Dorf *et al*, 2000; Fong *et al*, 1998; Goda *et al*, 2000; Harrison *et al*, 1998; Imai *et al*, 1997; Tong *et al*, 2000). The monocytes, when recruited to the site of injury, could secrete other chemokines that recruit additional leukocytes to the site of tissue injury and induce inflammation. Indeed, the FKN-CX3CR1 pair may participate in the generation and progression of inflammatory disorders within the brain and periphery. FKN has already been shown to play a role in a variety of pathological conditions related to inflammation, including atherosclerosis (Alexander, 2001; Greaves and Gordon, 2001; Greaves *et al*, 2001; McDermott *et al*, 2001), renal inflammation (Cockwell *et al*, 2002; Feng *et al*, 1999; Furuichi *et al*, 2001), airway inflammation (Fujimoto *et al*, 2001), psoriasis (Raychaudhuri *et al*, 2001), arthritis (Ruth *et al*, 2001; Volin *et al*, 2001), cardiac allograft rejection (Haskell *et al*, 2001), progression of acquired immunodeficiency syndrome (AIDS) (Faure *et al*, 2000; Foussat *et al*, 2001), and CNS inflammation (Boehme *et al*, 2000; Harrison *et al*, 1998; Hughes *et al*, 2002; Maciejewski-Lenoir *et al*, 1999; Nishiyori *et al*, 1998; Schwaeble *et al*, 1998; Zujovic *et al*, 2000). These observations suggest that FKN and its receptor have a unique role in regulation of the host response to disease (Fong *et al*, 1998; Harrison *et al*, 1998).

The balance of data demonstrates overwhelming support for FKN mediated recruitment of leukocytes. However, studies with FKN and CX3CR1 knockout mice (Cook *et al*, 2001; Jung *et al*, 2000) have tempered the importance of FKN and CX3CR1 in leukocyte recruitment. In studies (Cook *et al*, 2001) with FKN-deficient mice, responses to a variety of inflammatory stimuli were indistinguishable from those of wild-type mice in an intestine inflammation model system (Cook *et al*, 2001). In other reports (Jung *et al*, 2000), the absence of CX3CR1 did not interfere with either monocyte extravasation or dendritic cell migration and differentiation in a peritonitis model. Further, CX3CR1-deficient microglia exhibited proficient responses to peripheral nerve injury, indicating unimpaired neuronal-glial cross-talk in the absence of CX3CR1 (Jung *et al*, 2000). These findings suggest that other means of neuronal-glial linkage exist. Nevertheless, to elucidate the exact role of FKN and CX3CR1-expressing cells in disease pathogenesis, further studies are certainly required. Moreover, FKN-CX3CR1 interactions may mediate other inflammatory responses besides the recruitment of leukocytes. The following section will examine how FKN mediates communication between neurons and MP during homeostasis and disease.

FKN expression, structure, and regulation

FKN (neurotactin, CX3CL1) is a 373-amino acid, multidomain molecule found in a wide variety of tissues, including liver, intestine, kidney, and brain.




Figure 1 A proposed pathophysiological mechanism for how MP activation influences neuronal injury in HAD. During monocyte/macrophage maturation, macrophages acquire the ability to sustain productive HIV-1 infection. Release of progeny virion leads to infection of resident brain microglia. Uninfected or virus-infected MP can be immune-activated by a process that remains incompletely understood, but likely involves cytokines, chemokines, and cell-to-cell interactions. Released from injured neurons, the neuronal chemokine fractalkine (FKN) represents one pathway through which MP activation may occur. After recruiting MP to the site of injury, FKN may activate MP to produce neurotrophic/toxic factors that affect neuronal survival and induce CNS inflammation. In turn, recruited MP may also become infected and activation leading to the production of chemokines, cytokines, and glutamate. Chemokines, gp120, and whole virions may also interact with neuronal receptors to alter intracellular signal transduction pathways, leading to neuronal dysfunction and death.

Structural components of FKN include a 76–amino acid chemokine domain (CD) at the N-terminus, which is important in the binding, adhesion, and activation of its target cells (Harrison *et al*, 2001; Mizoue *et al*, 1999, 2001; Goda *et al*, 2000; Haskell *et al*, 2000). In addition, FKN has a 241–amino acid mucin-like stalk, which extends the chemokine domain away from the cell surface in order to aid in the adherence of CX3CR1-expressing cells (Fong *et al*, 2000). FKN also has an 18–amino acid stretch of hydrophobic residues that spans the cell membrane, and an extended C-terminus that anchors it to the cell surface (Cook *et al*, 2001; Hoover *et al*, 2000; Lucas *et al*, 2001). These unique structural features enable FKN to mediate chemotaxis, adherence, and activation of CX3CR1-expressing cells.

FKN is novel in that it is the only chemokine known to be expressed at higher levels within the CNS than in the periphery (Bazan *et al*, 1997). In

the CNS, FKN is constitutively expressed by neurons (Harrison *et al*, 1998; Hughes *et al*, 2002) (Figure 2) and can be induced by astrocytes (Hughes *et al*, 2002; Pereira *et al*, 2001; Zheng *et al*, 2002). It is up-regulated and released in response to proinflammatory stimuli, such as lipopolysaccharide (LPS), IL-1 β , tumor necrosis factor (TNF)- α , CD40L, and interferon (IFN)- γ (Fratelli *et al*, 2001; Fujimoto *et al*, 2001; Garcia *et al*, 2000; Hughes *et al*, 2002; Imaizumi *et al*, 2000; Pereira *et al*, 2001; Yoshida *et al*, 2001; Zheng *et al*, 2002). This up-regulation is believed to occur through activation of nuclear factor (NF)- κ B (Garcia *et al*, 2000).

FKN is also distinct from other chemokines, because it exists in both membrane-bound and soluble isoforms (Fong *et al*, 2000; Harrison *et al*, 2001; Mizoue *et al*, 2001). In response to excitotoxic stimuli, the membrane-spanning domain is rapidly cleaved and a soluble form of FKN is released from




Figure 2 FKN expression in human neuronal cells. Panel (A) shows mixed human cortical cells in culture that were stained for FKN (green) and panel (B) shows neurons double-stained for FKN and MAP-2 (yellow, neuronal marker) (200 \times). Results are representative of three independent experiments.

the neuronal cell surface (Chapman *et al*, 2000a; Zheng *et al*, 2000). Proteolytic cleavage of FKN is proposed to occur at a di-arginine sequence located next to the transmembrane domain (Bazan *et al*, 1997; Cook *et al*, 2001; Fong *et al*, 2000; Harrison *et al*, 1998, 2001). However, the exact location of the cleavage site remains to be confirmed (Tsou *et al*, 2001). FKN cleavage can be mediated by two distinct metalloproteinase-dependent activities: a constitutive FKN sheddase, which is active under normal cell culture conditions, and an inducible FKN sheddase that can be rapidly activated by phorbol esters, such as phorbol 12-myristate 13-acetate (PMA) (Garton *et al*, 2001; Tsou *et al*, 2001). Recently, inducible cleavage has been shown to be mediated by the TNF- α -converting enzyme (TACE), which belongs to a family of proteins containing a metalloprotease domain (Garton *et al*, 2001; Tsou *et al*, 2001).

The receptor for FKN, CX3CR1 (Combadiere *et al*, 1998), is expressed on monocytes (Cambien *et al*, 2001; Chapman *et al*, 2000b), dendritic cells (Dichmann *et al*, 2001), T lymphocytes (Fong *et al*, 1998; Foussat *et al*, 2000; Fraticelli *et al*, 2001), natural killer cells (Fong *et al*, 1998; Imai *et al*, 1997; Inngjerdingen *et al*, 2001), astrocytes (Dorf *et al*, 2000), neurons (Hughes *et al*, 2002; Meucci *et al*, 2000; Tong *et al*, 2000), and brain microglia (Boehme *et al*, 2000; Chapman *et al*, 2000a; Harrison *et al*, 1998; Hughes *et al*, 2002). Like other chemokine receptors, CX3CR1 (previously called V28) belongs to a family of GPCRs, which feature a seven-transmembrane domain, an extracellular N-terminus, and a cytoplasmic C-terminus. GPCRs interact with and signal through heterotrimeric guanine nucleotide-binding regulatory proteins (G-proteins).

Upon stimulation by a ligand, GPCRs undergo a conformational change that leads to activation of the G-protein by GDP-GTP exchange, followed by uncoupling of the G-protein from the receptor. Upon activation, G-proteins trigger a cascade of signaling events that regulate various cellular functions (Devi, 2000).

FKN functions: Cell adhesion and neuroprotection

In the brain, FKN is believed to regulate a complex network of paracrine and autocrine interactions between neurons and surrounding MP (Boehme *et al*, 2000; Harrison *et al*, 1998; Maciejewski-Lenoir *et al*, 1999), primarily through chemoattraction and adhesion. Both the soluble and membrane-bound forms of FKN are potent inducers of chemotaxis (Harrison *et al*, 2001). However, it is the membrane-bound form that enables FKN to immobilize CX3CR1-expressing cells, such as leukocytes (Boehme *et al*, 2000; Chapman *et al*, 2000a, 2000b; Combadiere *et al*, 1998; Dorf *et al*, 2000; Fong *et al*, 1998; Harrison *et al*, 1998; Imai *et al*, 1997; Tong *et al*, 2000). Mutation analyses and knockout mouse experiments have shown that specific residues within the FKN CD, such as Lys-7 and Arg-47, are important determinants in mediating binding, signaling, and adhesion of CX3CR1-expressing cells (Goda *et al*, 2000; Harrison *et al*, 2001; Haskell *et al*, 2000; Mizoue *et al*, 1999, 2001). Further, the adherence of FKN to CX3CR1-expressing leukocytes is believed to be integrin independent (Fong *et al*, 1998). Other studies suggest that adhesion of CX3CR1-expressing leukocytes is independent of G-protein activation (Haskell *et al*, 1999). It is

possible that the mucin-like domain of FKN may aid in adherence of CX3CR1-expression cells by extending the chemokine domain away from the cell surface in order to present it to trafficking leukocytes (Fong *et al*, 2000). Additionally, it is possible that the ability of FKN to mediate adhesion of trafficking cells may be a function of its slow receptor off-rate (Haskell *et al*, 2000). Nevertheless, it is clear that FKN and CX3CR1 fulfill important roles in leukocyte trafficking.

In addition to chemoattraction and adhesion, FKN may serve other functions, such as inhibition of HIV-1 infection and neuroprotection (Fong *et al*, 1998; Harrison *et al*, 1998; Haskell *et al*, 2000; Inngjerdingen *et al*, 2001; Tong *et al*, 2000). For example, FKN inhibits HIV-1 entry into CX3CR1-expressing cells (Faure *et al*, 2000) and inhibits neuronal injury induced by gp120 (Meucci *et al*, 2000), platelet-activating factor (PAF), and the regulatory HIV-1 gene product, Tat (Tong *et al*, 2000). FKN has also been shown to inhibit Fas-mediated death in microglia (Boehme *et al*, 2000). The protective functions of FKN are believed to be mediated through activation of signaling pathways involving the protein kinase, Akt (protein kinase B), and NF- κ B, which are major components of prosurvival signaling pathways in neurons and microglia (Boehme *et al*, 2000; Meucci *et al*, 2000).

FKN and macrophage activation: Dysregulation of neurotrophic/toxic factors?

Because CX3CR1 is highly expressed on MP, it is possible that FKN-CX3CR1 interactions play an important role in mediating MP immune activation. Upon binding to CX3CR1, FKN has been shown to stimulate TNF- α and IL-8 production in MP (Figure 3) (Zheng *et al*, 2002; Zujovic *et al*, 2000). Although many of the individual factors secreted by FKN-activated MP remain to be determined, it is known that HIV-1-infected and immune-activated MP are capable of producing a wide variety of toxic factors. These factors include proinflammatory cytokines such as




Figure 3 IL-8 (A) and TNF- α (B) production in cell supernatants from FKN-treated human monocyte-derived macrophages (MDM). After 14 days in culture, elutriated and recombinant human macrophage colony-stimulating factor (MCSF)-differentiated human MDM were treated with different concentrations of soluble FKN for 4 h. * $P < .01$ as compared to control. Results are expressed as an average \pm SD and are representative of three independent experiments.

TNF- α , IL-1 β (Sebire *et al*, 1993), glutamate (Jiang *et al*, 2001), arachidonic acid and its metabolites (Genis *et al*, 1992), PAF (Gelbard *et al*, 1994), quinolinic acid (Heyes *et al*, 1991; Kerr *et al*, 1998), NTox (Giulian *et al*, 1996), nitric oxide (NO) (Adamson *et al*, 1996), and reactive oxygen species (ROS) (Mollace *et al*, 2001). Alternatively, viral infection and FKN-mediated activation of MP may regulate production of trophic factors that mediate neuronal growth and repair (Lopez *et al*, 2001). A number of neurotrophic factors are secreted by MP (Barnea *et al*, 1996; Elkabes *et al*, 1996), including brain-derived neurotrophic factor (BDNF) (Kerschensteiner *et al*, 1999; Miwa *et al*, 1997), β -nerve growth factor (β NGF) (Caroleo *et al*, 2001; Garaci *et al*, 1999; Lopez *et al*, 2001), transforming growth factor-beta (TGF- β) (Chao *et al*, 1995), neurotrophin-3 (NT3) (Kullander *et al*, 1997; Loy *et al*, 1994; Mallat *et al*, 1989; Rocamora *et al*, 1996; Saad *et al*, 1991), and glial-derived neurotrophic factor (GDNF) (Batchelor *et al*, 1999). Withdrawal or dysregulation of these factors can result in neuronal injury and death (Deshmukh *et al*, 1996). Through enhanced neurotoxin secretion and dysregulated neurotrophin production, HIV-1-infected and FKN-activated MP may induce neuronal injury and death in HIVE (Aquaro *et al*, 2000; Conant *et al*, 1998; Cotter *et al*, 1999a; Fischer-Smith *et al*, 2001; Gabuzda *et al*, 1998; Gendelman, 1997; Glass *et al*, 1995; Koenig *et al*, 1986; Lopez *et al*, 2001; Nath and Geiger, 1998; Perno *et al*, 1997; Strizki *et al*, 1996; Wiley *et al*, 1986; Zheng and Gendelman, 1997).

FKN-induced secretory factor production is believed to occur through activation of intracellular signaling pathways (Cambien *et al*, 2001; Zheng *et al*, 2002). Therefore, the following section will discuss the relevant intracellular signaling pathways resulting from FKN-mediated activation of CX3CR1 expressing-MP.

FKN-mediated signal transduction pathways

Binding of FKN to CX3CR1 on MP initiates multiple signal transduction pathways and leads to the activation of a wide variety of protein kinases, including the tyrosine kinases (the Src tyrosine kinase family and Syk tyrosine kinase family), calcium calmodulin kinase (CaMK), protein kinase C (PKC), phosphatidylinositol 3-kinase (PI 3-kinase), protein kinase B, mitogen-activated protein kinases (MAP kinases), and NF- κ B (Cambien *et al*, 2001; Garcia *et al*, 2000). Activation of these signal transduction pathways leads to elevation of cytosolic free calcium and modifications in enzymes, ion channels, transcriptional activators, and transcriptional regulators (Cambien *et al*, 2001; Iismaa *et al*, 1995).

Several studies have shown that binding of FKN to CX3CR1 induces the activation of MAP kinases (Figure 4) (Cambien *et al*, 2001; Zheng *et al*, 2002). Activation of MAP kinase pathways stimulate




Figure 4 Overview of proposed MAP kinase and protein kinases B and C signal transduction events in mononuclear phagocyte (MP)-mediated production of proinflammatory factors or neurotrophic/toxic factors. FKN can bind the chemokine receptor CX3CR1 on MP and activate MAP kinase signaling through the α or $\beta\gamma$ subunit of G-protein, which can further activate one or more types of MAP kinases. These MAP kinases include extracellular signal-related kinases (ERK1 and ERK2) and stress-activated protein kinases (SAPK1/JNK1 and SAPK2/p38). In addition, FKN can also activate intracellular signaling pathways, such as increasing cytosolic free calcium, activation of phosphatidylinositol 3-kinase (PI-3K) and protein kinase B (PKB), and alteration of protein kinase C, which further activate MP. This activation causes the production of proinflammatory factors or multiple neurotrophic/toxic factors. The inhibitors for different kinase pathways can be used as tools to elucidate the signaling pathways involved in MP activation events. Some of the stimulation pathways may increase the cellular activation state and cause overproduction of cytokines or neurotoxins, which mediate MP-induced neuronal injury in HAD.

cell growth and differentiation by regulating gene translation and expression (Lopez-Illasaca, 1998; Lopez-Illasaca *et al*, 1997). There are three distinct MAP kinase cascades (Figure 4): c-Jun N-terminal kinase/stress-activated protein kinase (JNK/SAPK), p38 (Lopez-Illasaca, 1998), and extracellular signal-related kinases (ERK1/ERK2). The JNK/SAPK pathway is induced by exposure to ultraviolet radiation, heat shock, or inflammatory cytokines. The p38 pathway is activated in response to inflammatory cytokines, endotoxins, and osmotic stress. The ERK pathway is stimulated following binding of extracellular growth factors (for example, epidermal growth factor, EGF) to tyrosine kinase-linked receptors. It appears that each of the three MAP kinase pathways, ERK1/2, p38, and JNK, are activated during the binding of FKN to CX3CR1 on monocyte-derived macrophages (MDM) (Zheng *et al*, 2002). Interestingly, production of MIP-1 β and IL-8 by FKN-activated MP can be blocked by MAP kinase inhibitors, such as PD98056 (ERK), SB203850 (p38), and SP600125 (JNK) (Zheng *et al*, 2002). Activation of MAP kinase signaling appears to be critical for FKN-induced capture, adhesion, and activation of MP (Cambien *et al*, 2001; Kansra *et al*, 2001; Zheng *et al*, 2002). Thus, multiple protein kinases appear to be involved in mediating the effects of FKN upon

its target cells. The elucidation of specific pathways through which FKN induces MP activation and regulates production of neurotrophic/toxic factors will be critical to understanding disease pathogenesis in HAD and other neurodegenerative disorders.

Summary

In summary, the evidence presented within this review suggests that the expression and production of neuronal chemokines, such as FKN, may be a compensatory or reparative response to injury in the brain. As such, we propose that in HAD, viral proteins, such as gp120, induce neuronal injury, leading to the up-regulation or release of FKN. Acting as a "distress signal," FKN may then recruit and activate CX3CR1-expressing MP to the site of injury (Fong *et al*, 2000; Harrison *et al*, 2001). Dysregulation of this response may result in further neuronal injury as MP themselves become activated to produce neurotoxins. This, in turn, may induce a cycle of inflammation and injury. Identification of the pathways through which FKN induces MP activation could lead to the development of agents that impede or prevent further neuronal injury in HAD and other neurodegenerative disorders.

References

- Adamson DC, Wildemann B, Sasaki M, Glass JD, McArthur JC, Christov VI, Dawson TM, Dawson VL (1996). Immunologic NO synthase: elevation in severe AIDS dementia and induction by HIV-1 gp41. *Science* **274**: 1917–1926.
- Albright AV, Shieh JT, Itoh T, Lee B, Pleasure D, O'connor MJ, Doms RW, Gonzalez-Scarano F (1999). Microglia express CCR5, CXCR4, and CCR3, but of these, CCR5 is the principal coreceptor for human immunodeficiency virus type 1 dementia isolates. *J Virol* **73**: 205–213.
- Alexander RW (2001). Cytokine receptor CX3CR-1 and fractalkine: new factors in the atherosclerosis drama? *Circ Res* **89**: 376–377.
- Aquaro S, Panti S, Caroleo MC, Balestra E, Cenci A, Forbici F, Ippolito G, Mastino A, Testi R, Mollace V, *et al* (2000). Primary macrophages infected by human immunodeficiency virus trigger CD95-mediated apoptosis of uninfected astrocytes. *J Leukoc Biol* **68**: 429–435.
- Asare E, Dunn G, Glass J, McArthur J, Luthert P, Lantos P, Everall I (1996). Neuronal pattern correlates with the severity of human immunodeficiency virus-associated dementia complex. *Am J Pathol* **148**: 31–38.
- Baggiolini M, Dewald B, Moser B (1997). Human chemokines: an update. *Annu Rev Immunol* **15**: 675–705.
- Barnea A, Aguila-Mansilla N, Chute HT, Welcher AA (1996). Comparison of neurotrophin regulation of human and rat neuropeptide Y (NPY) neurons: induction of NPY production in aggregate cultures derived from rat but not from human fetal brains. *Brain Res* **732**: 52–60.
- Batchelor PE, Liberatore GT, Wong JY, Porritt MJ, Frerichs F, Donnan GA, Howells DW (1999). Activated macrophages and microglia induce dopaminergic sprouting in the injured striatum and express brain-derived neurotrophic factor and glial cell line-derived neurotrophic factor. *J Neurosci* **19**: 1708–1716.
- Bazan J, Bacon K, Hardiman G, Wang W, Soo K, Rossi D, Greaves D, Zlotnik A, Schall T (1997). A new class of membrane-bound chemokine with a CX3C motif. *Nature* **385**: 640–644.
- Biber K, Sauter A, Brouwer N, Copray SC, Boddeke HW (2001). Ischemia-induced neuronal expression of the microglia attracting chemokine secondary lymphoid-tissue chemokine (SLC). *Glia* **34**: 121–133.
- Boehme SA, Lio FM, Maciejewski-Lenoir D, Bacon KB, Conlon PJ (2000). The chemokine fractalkine inhibits Fas-mediated cell death of brain microglia. *J Immunol* **165**: 397–403.
- Broder CC, Collman RG (1997). Chemokine receptors and HIV. *J Leukoc Biol* **62**: 20–29.
- Cambien B, Pomeranz M, Schmid-Antomarchi H, Millet MA, Breittmayer V, Rossi B, Schmid-Alliana A (2001). Signal transduction pathways involved in soluble fractalkine-induced monocytic cell adhesion. *Blood* **97**: 2031–2037.
- Caroleo MC, Costa N, Bracci-Laudiero L, Aloe L (2001). Human monocyte/macrophages activated by exposure to LPS overexpress NGF and NGF receptors. *J Neuroimmunol* **113**: 193–201.
- Carpenter CC, Cooper DA, Fischl MA, Gatell JM, Gazzard BG, Hammer SM, Hirsch MS, Jacobsen DM,

- Katzenstein DA, Montaner JS, et al (2000). Antiretroviral therapy in adults: updated recommendations of the International AIDS Society–USA Panel. *JAMA* **283**: 381–390.
- Chao CC, Hu S, Sheng WS, Tsang M, Peterson PK (1995). Tumor necrosis factor- α mediates the release of bioactive transforming growth factor β in murine microglial cell cultures. *Clin Immunol Immunopathol* **77**: 358–365.
- Chapman GA, Moores K, Harrison D, Campbell CA, Stewart BR, Strijbos PJ (2000a). Fractalkine cleavage from neuronal membranes represents an acute event in the inflammatory response to excitotoxic brain damage. *J Neurosci (Online)* **20**: RC87.
- Chapman GA, Moores KE, Gohil J, Berkhout TA, Patel L, Green P, Macphee CH, Stewart BR (2000b). The role of fractalkine in the recruitment of monocytes to the endothelium. *Eur J Pharmacol* **392**: 189–195.
- Cockwell P, Chakravorty SJ, Girdlestone J, Savage CO (2002). Fractalkine expression in human renal inflammation. *J Pathol* **196**: 85–90.
- Combadiere C, Salzwedel K, Smith ED, Tiffany HL, Berger EA, Murphy PM (1998). Identification of CX3CR1. A chemotactic receptor for the human CX3C chemokine fractalkine and a fusion coreceptor for HIV-1. *J Biol Chem* **273**: 23799–23804.
- Conant K, Garzino-Demo A, Nath A, McArthur JC, Halliday W, Power C, Gallo RC, Major EO (1998). Induction of monocyte chemoattractant protein-1 in HIV-1 Tat-stimulated astrocytes and elevation in AIDS dementia. *Proc Natl Acad Sci USA* **95**: 3117–3121.
- Cook DN, Chen SC, Sullivan LM, Manfra DJ, Wiekowski MT, Prosser DM, Vassileva G, Lira SA (2001). Generation and analysis of mice lacking the chemokine fractalkine. *Mol Cell Biol* **21**: 3159–3165.
- Cotter R, Zheng J, Gendelman HE (1999a). The role of mononuclear phagocytes in neurodegenerative disorders: lessons from multiple sclerosis, Alzheimer's disease and HIV-1 dementia. In: *Advances in neurodegenerative disorders*. Marwah J, Teitelbaum H (eds). Prominent Press: Scottsdale, AZ, pp 203–241.
- Cotter R, Zheng J, Niemann D, Thomas E, Gendelman H (1999b). CD40L activation of mononuclear phagocytes: regulation of HIV-1 replication and beta-chemokine production. Paper presented at XIth International Congress of Virology, Interbational Union of Microbiological Societies, Sydney, Australia, August 9–13.
- Cotter RL, Zheng J, Che M, Niemann D, Liu Y, He J, Thomas E, Gendelman HE (2001). Regulation of human immunodeficiency virus type 1 infection, beta chemokine production, and CCR5 expression in CD40L-stimulated macrophages: immune control of viral entry. *J Virol* **75**: 4308–4320.
- Coughlan CM, McManus CM, Sharron M, Gao Z, Murphy D, Jaffer S, Choe W, Chen W, Hesselgesser J, Gaylord H, et al (2000). Expression of multiple functional chemokine receptors and monocyte chemoattractant protein-1 in human neurons [In Process Citation]. *Neuroscience* **97**: 591–600.
- Desbaillets I, Tada M, de Tribolet N, Diserens AC, Hamou MF, Van Meir EG (1994). Human astrocytomas and glioblastomas express monocyte chemoattractant protein-1 (MCP-1) in vivo and in vitro. *Int J Cancer* **58**: 240–247.
- Deshmukh M, Vasilakos J, Deckwerth TL (1996). Genetic and metabolic status of NGF-deprived sympathetic neurons saved by an inhibitor of ICE family proteases. *J Cell Biol* **135**: 1341–1354.
- Devi LA (2000). G-protein-coupled receptor dimers in the lime light. *Trends Pharmacol Sci* **21**: 324–326.
- Dichmann S, Herouy Y, Purlis D, Rheinen H, Gebicke-Harter P, Norgauer J (2001). Fractalkine induces chemotaxis and actin polymerization in human dendritic cells. *Inflamm Res* **50**: 529–533.
- Dickson D, Lee S, Hatch W, Mattiace L, Brosnan C, Lyman W (1994). Macrophages and microglia in HIV-related CNS neuropathology. In: *HIV, AIDS, and the brain*. Price R, Perry S (eds). Raven: New York, pp 99–118.
- Dorf ME, Berman MA, Tanabe S, Heesen M, Luo Y (2000). Astrocytes express functional chemokine receptors. *J Neuroimmunol* **111**: 109–121.
- Dragic T, Litwin V, Allaway GP, Martin SR, Huang Y, Nagashima KA, Cayanan C, Maddon PJ, Koup RA, Moore JP, Paxton WA (1996). HIV-1 entry into CD4+ cells is mediated by the chemokines receptor CC-CKR-5. *Nature* **381**: 667–673.
- Elkabes S, DiCicco-Bloom EM, Black IB (1996). Brain microglia/macrophages express neurotrophins that selectively regulate microglial proliferation and function. *J Neurosci* **16**: 2508–2521.
- Endres MJ, Clapham PR, Marsh M, Ahuja M, Turner JD, McKnight A, Thomas JF, Stoebenau-Haggarty B, Choe S, Vance PJ, et al (1996). CD4-independent infection by HIV-2 is mediated by fusin/CXCR4. *Cell* **87**: 745–756.
- Faure S, Meyer L, Costagliola D, Vaneensberghe C, Genin E, Autran B, Delfraissy JF, McDermott DH, Murphy PM, Debre P, et al (2000). Rapid progression to AIDS in HIV+ individuals with a structural variant of the chemokine receptor CX3CR1. *Science* **287**: 2274–2277.
- Feng L, Chen S, Garcia GE, Xia Y, Siani MA, Botti P, Wilson CB, Harrison JK, Bacon KB (1999). Prevention of crescentic glomerulonephritis by immunoneutralization of the fractalkine receptor CX3CR1 rapid communication. *Kidney Int* **56**: 612–620.
- Fischer-Smith T, Croul S, Sverstiuk AE, Capini C, L'Heureux D, Regulier EG, Richardson MW, Amini S, Morgello S, Khalili K, Rappaport J (2001). CNS invasion by CD14+/CD16+ peripheral blood-derived monocytes in HIV dementia: perivascular accumulation and reservoir of HIV infection. *J NeuroVirol* **7**: 528–541.
- Fong AM, Erickson HP, Zachariah JP, Poon S, Schamberg NJ, Imai T, Patel DD (2000). Ultrastructure and function of the fractalkine mucin domain in CX(3)C chemokine domain presentation. *J Biol Chem* **275**: 3781–3786.
- Fong AM, Robinson LA, Steeber DA, Tedder TF, Yoshie O, Imai T, Patel DD (1998). Fractalkine and CX3CR1 mediate a novel mechanism of leukocyte capture, firm adhesion, and activation under physiologic flow. *J Exp Med* **188**: 1413–1419.
- Foussat A, Bouchet-Delbos L, Berrebi D, Durand-Gasselini I, Coulomb-L'Hermine A, Krzysiek R, Galanaud P, Levy Y, Emilie D (2001). Deregulation of the expression of the fractalkine/fractalkine receptor complex in HIV-1-infected patients. *Blood* **98**: 1678–1686.
- Foussat A, Coulomb-L'Hermine A, Gosling J, Krzysiek R, Durand-Gasselini I, Schall T, Balian A, Richard Y, Galanaud P, Emilie D (2000). Fractalkine receptor expression by T lymphocyte subpopulations and in vivo production of fractalkine in human. *Eur J Immunol* **30**: 87–97.

- Fraticelli P, Sironi M, Bianchi G, D'Ambrosio D, Albanesi C, Stoppacciaro A, Chieppa M, Allavena P, Ruco L, Girolomoni G, et al (2001). Fractalkine (CX3CL1) as an amplification circuit of polarized Th1 responses. *J Clin Invest* **107**: 1173–1181.
- Fujimoto K, Imaizumi T, Yoshida H, Takanashi S, Okumura K, Satoh K (2001). Interferon-gamma stimulates fractalkine expression in human bronchial epithelial cells and regulates mononuclear cell adherence. *Am J Respir Cell Mol Biol* **25**: 233–238.
- Furuichi K, Wada T, Iwata Y, Sakai N, Yoshimoto K, Shimizu M, Kobayashi K, Takasawa K, Kida H, Takeda S, et al (2001). Upregulation of fractalkine in human crescentic glomerulonephritis. *Nephron* **87**: 314–320.
- Gabuzda D, He J, Ohagen A, Vallat A (1998). Chemokine receptors in HIV-1 infection of the central nervous system. *Immunology* **10**: 203–213.
- Gabuzda D, Wang J (1999). Chemokine receptors and virus entry in the central nervous system. *J NeuroVirol* **5**: 643–658.
- Gabuzda D, Wang J (2000). Chemokine receptors and mechanisms of cell death in HIV neuropathogenesis. *J NeuroVirol* **6 (Suppl 1)**: S24–S32.
- Gabuzda D, Wang J, Gorry P (2002). HIV-1-associated dementia. In: *Chemokines and the nervous system*. Ransohoff RM, Suzuki K, Proudfoot AEI, Hickey WF, Harrison JK (eds). Elsevier Science: Amsterdam, pp 345–360.
- Garaci E, Caroleo MC, Aloe L, Aquaro S, Piacentini M, Costa N, Amendola A, Micera A, Calio R, Perno CF, Levi-Montalcini R (1999). Nerve growth factor is an autocrine factor essential for the survival of macrophages infected with HIV. *Proc Natl Acad Sci USA* **96**: 14013–14018.
- Garcia GE, Xia Y, Chen S, Wang Y, Ye RD, Harrison JK, Bacon KB, Zerwes HG, Feng L (2000). NF-kappaB-dependent fractalkine induction in rat aortic endothelial cells stimulated by IL-1beta, TNF-alpha, and LPS. *J Leukoc Biol* **67**: 577–584.
- Garton KJ, Gough PJ, Blobel CP, Murphy G, Greaves DR, Dempsey PJ, Raines EW (2001). Tumor necrosis factor-alpha-converting enzyme (ADAM17) mediates the cleavage and shedding of fractalkine (CX3CL1). *J Biol Chem* **276**: 37993–38001.
- Gelbard H, Nottet H, Dzenko K, Jett M, Genis P, White R, Wang L, Choi Y-B, Zhang D, Lipton S, et al (1994). Platelet-activating factor: a candidate human immunodeficiency virus type-1 infection neurotoxin. *J Virol* **68**: 4628–4635.
- Gelbard HA, Epstein LG (1995). HIV-1 encephalopathy in children. *Curr Opin Pediatr* **7**: 655–662.
- Gendelman H, Persidsky Y, Ghorpade A, Limoges J, Stins M, Fiala M, Morrisett R (1997). The neuropathogenesis of AIDS dementia complex. *AIDS* **11**: S35–S45.
- Gendelman HE (1997). The neuropathogenesis of HIV-1 dementia. In: *The neurology of AIDS*. Gendelman HE, Lipton SA, Epstein LG, Swindells S (eds). Chapman and Hall: New York, pp 1–10.
- Genis P, Jett M, Bernton E, Boyle T, Gelbard H, Dzenko K, Keane R, Resnick L, Mizrachi Y, Volsky D, et al (1992). Cytokines and arachidonic metabolites produced during human immunodeficiency virus (HIV)-infected macrophage-astroglia interactions: implications for the neuropathogenesis of HIV disease. *J Exp Med* **176**: 1703–1718.
- Ghorpade A, Xia MQ, Hyman BT, Persidsky Y, Nukuna A, Bock P, Che M, Limoges J, Gendelman HE, Mackay CR (1998). Role of the beta-chemokine receptors CCR3 and CCR5 in human immunodeficiency virus type 1 infection of monocytes and microglia. *J Virol* **72**: 3351–3361.
- Giulian D, Noonan C, Vaca K (1990). HIV-1 infected mononuclear phagocytes release neurotoxins. *Science* **250**: 1593–1595.
- Giulian D, Yu J, Xia L, Tom D, Li J, Lin SN, Schwarz R, Noonan C (1996). Study of receptor-mediated neurotoxins released by HIV-1 infected mononuclear phagocytes found in human brain. *J Neurosci* **16**: 3139–3153.
- Glass JD, Fedor H, Wesselingh SL, McArthur JC (1995). Immunocytochemical quantitation of human immunodeficiency virus in the brain: correlations with dementia. *Ann Neurol* **38**: 755–762.
- Gleichmann M, Gillen C, Czardybon M, Bosse F, Greiner-Petter R, Auer J, Muller HW (2000). Cloning and characterization of SDF-1gamma, a novel SDF-1 chemokine transcript with developmentally regulated expression in the nervous system. *Eur J Neurosci* **12**: 1857–1866.
- Goda S, Imai T, Yoshie O, Yoneda O, Inoue H, Nagano Y, Okazaki T, Imai H, Bloom ET, Domae N, Umehara H (2000). CX3C-chemokine, fractalkine-enhanced adhesion of THP-1 cells to endothelial cells through integrin-dependent and -independent mechanisms. *J Immunol* **164**: 4313–4320.
- Greaves DR, Hakkinen T, Lucas AD, Liddiard K, Jones E, Quinn CM, Senaratne J, Green FR, Tyson K, Boyle J, et al (2001). Linked chromosome 16q13 chemokines, macrophage-derived chemokine, fractalkine, and thymus- and activation-regulated chemokine are expressed in human atherosclerotic lesions. *Arterioscler Thromb Vasc Biol* **21**: 923–929.
- Greaves DR, Gordon S (2001). Immunity, atherosclerosis and cardiovascular disease. *Trends Immunol* **22**: 180–181.
- Harrison JK, Fong AM, Swain PA, Chen S, Yu YR, Salafranca MN, Greenleaf WB, Imai T, Patel DD (2001). Mutational analysis of the fractalkine chemokine domain: basic amino acid residues differentially contribute to CX3CR1 binding, signaling, and cell adhesion. *J Biol Chem* **276**: 8.
- Harrison JK, Jiang Y, Chen S, Xia Y, Maciejewski D, McNamara RK, Streit WJ, Salafranca MN, Adhikari S, Thompson DA, et al (1998). Role for neuronally derived fractalkine in mediating interactions between neurons and CX3CR1-expressing microglia. *Proc Natl Acad Sci USA* **95**: 10896–10901.
- Haskell CA, Cleary MD, Charo IF (1999). Molecular uncoupling of fractalkine-mediated cell adhesion and signal transduction. Rapid flow arrest of CX3CR1-expressing cells is independent of G-protein activation. *J Biol Chem* **274**: 10053–10058.
- Haskell CA, Cleary MD, Charo IF (2000). Unique role of the chemokine domain of fractalkine in cell capture. Kinetics of receptor dissociation correlate with cell adhesion. *J Biol Chem* **275**: 34183–34189.
- Haskell CA, Hancock WW, Salant DJ, Gao W, Csizmadia V, Peters W, Faia K, Fituri O, Rottman JB, Charo IF (2001). Targeted deletion of CX(3)CR1 reveals a role for fractalkine in cardiac allograft rejection. *J Clin Invest* **108**: 679–688.
- He J, Chen Y, Farzan M, Choe H, Ohagen A, Gartner S, Busciglio J, Yang X, Hofmann W, Newman W,

- et al (1997). CCR3 and CCR5 are co-receptors for HIV-1 infection of microglia. *Nature* **385**: 645–649.
- Hesselgesser J, Horuk R (1999). Chemokine and chemokine receptor expression in the central nervous system. *J NeuroVirol* **5**: 13–26.
- Heyes MP, Brew BB, Martin A, Price RW, Salazar A, Sidtis JJ, Yergey JA, Mouradian MM, Sadler AE, Keilp J, et al (1991). Quinolinic acid in cerebrospinal fluid and serum in HIV-1 infection: relationship to clinical and neurological status. *Ann Neurol* **29**: 202–209.
- Hoover DM, Mizoue LS, Handel TM, Lubkowski J (2000). The crystal structure of the chemokine domain of fractalkine shows a novel quaternary arrangement. *J Biol Chem* **275**: 23187–23193.
- Horuk R, Martin A, Hesselgesser J, Hadley T, Lu ZH, Wang ZX, Peiper SC (1996). The Duffy antigen receptor for chemokines: structural analysis and expression in the brain. *J Leukoc Biol* **59**: 29–38.
- Hughes PM, Botham MS, Frentzel S, Mir A, Perry VH (2002). Expression of fractalkine (CX3CL1) and its receptor, CX3CR1, during acute and chronic inflammation in the rodent CNS. *Glia* **37**: 314–327.
- Iismaa T, Biden T, Shine J (1995). G protein-coupled receptors. RG Landes: Austin, TX.
- Imai T, Hieshima K, Haskell C, Baba M, Nagira M, Nishimura M, Kakizaki M, Takagi S, Nomiyama H, Schall TJ, Yoshie O (1997). Identification and molecular characterization of fractalkine receptor CX3CR1, which mediates both leukocyte migration and adhesion. *Cell* **91**: 521–530.
- Imaizumi T, Matsumiya T, Fujimoto K, Okamoto K, Cui X, Ohtaki U, Hidemi Yoshida, Satoh K (2000). Interferon-gamma stimulates the expression of CX3CL1/fractalkine in cultured human endothelial cells. *Tohoku J Exp Med* **192**: 127–139.
- Inngjerdigen M, Damaj B, Maghazachi AA (2001). Expression and regulation of chemokine receptors in human natural killer cells. *Blood* **97**: 367–375.
- Janssen R, Cornblath D, Epstein L, Foa R, McArthur J, Price R, Asbury A, Beckett A, Benson D, Bridge T, et al (1991). Nomenclature and research case definitions for neurological manifestations of human immunodeficiency virus type 1 (HIV-1) infection. *Neurology* **41**: 778–785.
- Jiang Z, Piggee C, Heyes MP, Murphy C, Quearry B, Bauer M, Zheng J, Gendelman HE, Markey SP (2001). Glutamate is a mediator of neurotoxicity in secretions of activated HIV-1-infected macrophages. *J Neuroimmunol* **117**: 97–107.
- Jung S, Aliberti J, Graemmel P, Sunshine MJ, Kreutzberg GW, Sher A, Littman DR (2000). Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green fluorescent protein reporter gene insertion. *Mol Cell Biol* **20**: 4106–4114.
- Kansra V, Groves C, Gutierrez-Ramos JC, Polakiewicz RD (2001). Phosphatidylinositol 3-kinase-dependent extracellular calcium influx is essential for CX(3)CR1-mediated activation of the mitogen-activated protein kinase cascade. *J Biol Chem* **276**: 31831–31838.
- Karpus WJ (2001). Chemokines and central nervous system disorders. *J NeuroVirol* **7**: 493–500.
- Kaul M, Lipton SA (1999). Chemokines and activated macrophages in HIV gp120-induced neuronal apoptosis [In Process Citation]. *Proc Natl Acad Sci USA* **96**: 8212–8216.
- Kelder W, McArthur JC, Nance-Sproson T, McClernon D, Griffin DE (1998). β -Chemokines MCP-1 and RANTES are selectively increased in cerebrospinal fluid of patients with human immunodeficiency virus-associated dementia. *Ann Neurol* **44**: 831–835.
- Kerr SJ, Armati PJ, Guillemin GJ, Brew BJ (1998). Chronic exposure of human neurons to quinolinic acid results in neuronal changes consistent with AIDS dementia complex. *AIDS* **12**: 355–363.
- Kerschensteiner M, Gallmeier E, Behrens L, Leal VV, Misgeld T, Klinkert WE, Kolbeck R, Hoppe E, Oropeza-Wekerle RL, Bartke I, et al (1999). Activated human T cells, B cells, and monocytes produce brain-derived neurotrophic factor in vitro and in inflammatory brain lesions: a neuroprotective role of inflammation? *J Exp Med* **189**: 865–870.
- Kitai R, Zhao ML, Zhang N, Hua LL, Lee SC (2000). Role of MIP-1 β and RANTES in HIV-1 infection of microglia: inhibition of infection and induction by IFN β . *J Neuroimmunol* **110**: 230–239.
- Klein R, Williams K, Alvarez-Hernandez X, Westmoreland S, Force T, Lackner A, Luster A (1999). Chemokine receptor expression and signaling in macaque and human fetal neurons and astrocytes: implications for the neuropathogenesis of AIDS. *J Immunol* **163**: 1636–1646.
- Koenig S, Gendelman HE, Orenstein JM, Canto MCD, Pezeshkpour GH, Yungbluth M, Janotta F, Aksamit A, Martin MA, Fauci AS (1986). Detection of AIDS virus in macrophages in brain tissue from AIDS patients with encephalopathy. *Science* **233**: 1089–1093.
- Kornbluth RS, Kee K, Richman DD (1998). CD40 ligand (CD154) stimulation of macrophages to produce HIV-1-suppressive beta-chemokines. *Proc Natl Acad Sci USA* **95**: 5205–5210.
- Krebs FC, Ross H, McAllister J, Wigdahl B (2000). HIV-1-associated central nervous system dysfunction. *Adv Pharmacol* **49**: 315–385.
- Kullander K, Kylberg A, Ebendal T (1997). Specificity of neurotrophin-3 determined by loss-of-function mutagenesis. *J Neurosci Res* **50**: 496–503.
- Kutsch O, Oh J, Nath A, Benveniste EN (2000). Induction of the chemokines interleukin-8 and IP-10 by human immunodeficiency virus type 1 tat in astrocytes. *J Virol* **74**: 9214–9221.
- Lavi E, Strizki JM, Ulrich AM, Zhang W, Fu L, Wang Q, O'Connor M, Hoxie JA, Gonzalez-Scarano F (1997). CXCR-4 (fusin), a co-receptor for the type 1 human immunodeficiency virus (HIV-1) is expressed in the human brain in a variety of cell types, including microglia and neurons. *Am J Pathol* **151**: 1035–1042.
- Letendre SL, Lanier ER, McCutchan JA (1999). Cerebrospinal fluid beta chemokine concentrations in neurocognitively impaired individuals infected with human immunodeficiency virus type 1. *J Infect Dis* **180**: 310–319.
- Lopez A, Bauer MA, Erichsen DA, Peng H, Gendelman L, Shibata A, Gendelman HE, Zheng J (2001). The regulation of neurotrophic factor activities following HIV-1 infection and immune activation of mononuclear phagocytes. Paper presented at the 31st Annual Meeting of the Society for Neuroscience, San Diego, California, November 10–15.
- Lopez-Illasaca M (1998). Signaling from G-protein coupled receptors to mitogen activated protein (MAP) kinase cascades. *Biochemical Pharmacol* **56**: 269–277.

- Lopez-Illasaca M, Crespo P, Pellici PG, Gutkind JS, Wetzker R (1997). Linkage of G protein-coupled receptors to the MAPK signaling pathway through Pl 3-kinase γ . *Science* **275**: 394–397.
- Loy R, Tagliabatella G, Angelucci L, Heyer D, Perez-Polo R (1994). Regional CNS uptake of blood-borne nerve growth factor. *J Neurosci Res* **39**: 339–346.
- Lucas AD, Chadwick N, Warren BF, Jewell DP, Gordon S, Powrie F, Greaves DR (2001). The transmembrane form of the CX3CL1 chemokine fractalkine is expressed predominantly by epithelial cells in vivo. *Am J Pathol* **158**: 855–866.
- Luster AD (1998). Chemokines-chemotactic cytokines that mediate inflammation. *N Engl J Med* **338**: 436–445.
- Ma M, Geiger JD, Nath A (1994). Characterization of a novel binding site for the human immunodeficiency virus type 1 envelope protein gp120 on human fetal astrocytes. *J Virol* **68**: 6824–6828.
- Maciejewski-Lenoir D, Chen S, Feng L, Maki R, Bacon KB (1999). Characterization of fractalkine in rat brain cells: migratory and activation signals for CX3CR1-expressing microglia. *J Immunol* **163**: 1628–1635.
- Mackay CR (1996). Chemokine receptors and T cell chemotaxis. *J Exp Med* **184**: 799–802.
- Mallat M, Houlgatte R, Brachet P, Prochiantz A (1989). Lipopolysaccharide-stimulated rat brain macrophages release NGF in vitro. *Dev Biol* **133**: 309–311.
- Marder K, Albert S, Dooneief G, Stern Y, Ramachandran G, Epstein L (1996). Clinical confirmation of the american academy of neurology algorithm for HIV-1-associated cognitive/motor disorder. *Neurology* **47**: 1247–1253.
- Masliah E (1996). In vivo modeling of HIV-1 mediated neurodegeneration. *Am J Pathol* **149**: 745–750.
- Masliah E, DeTeresa RM, Mallory ME, Hansen LA (2000). Changes in pathological findings at autopsy in AIDS cases for the last 15 years. *Aids* **14**: 69–74.
- McArthur JC, Sacktor N, Selnes O (1999). Human immunodeficiency virus-associated dementia. *Semin Neurol* **19**: 129–150.
- McDermott DH, Halcox JP, Schenke WH, Waclawiw MA, Merrell MN, Epstein N, Quyyumi AA, Murphy PM (2001). Association between polymorphism in the chemokine receptor CX3CR1 and coronary vascular endothelial dysfunction and atherosclerosis. *Circ Res* **89**: 401–407.
- Meucci O, Fatatis A, Simen AA, Bushell TJ, Gray PW, Miller RJ (1998). Chemokines regulate hippocampal neuronal signaling and gp120 neurotoxicity. *Proc Natl Acad Sci USA* **95**: 14500–14505.
- Meucci O, Fatatis A, Simen AA, Miller RJ (2000). Expression of CX3CR1 chemokine receptors on neurons and their role in neuronal survival. *Proc Natl Acad Sci USA* **97**: 8075–8080.
- Miller RJ, Meucci O (1999). AIDS and the brain: is there a chemokine connection? *Trends Neurosci* **22**: 471–479.
- Minami M, Satoh M (2000). Chemokines as mediators for intercellular communication in the brain (in Japanese). *Nippon Yakurigaku Zasshi* **115**: 193–200.
- Miwa T, Furukawa S, Nakajima K, Furukawa Y, Kohsaka S (1997). Lipopolysaccharide enhances synthesis of brain-derived neurotrophic factor in cultured rat microglia. *J Neurosci Res* **50**: 1023–1029.
- Mizoue LS, Bazan JF, Johnson EC, Handel TM (1999). Solution structure and dynamics of the CX3C chemokine domain of fractalkine and its interaction with an N-terminal fragment of CX3CR1. *Biochemistry* **38**: 1402–1414.
- Mizoue LS, Sullivan SK, King DS, Kledal TN, Schwartz TW, Bacon KB, Handel TM (2001). Molecular determinants of receptor binding and signaling by the CX3C chemokine fractalkine. *J Biol Chem* **276**: 33906–33914.
- Mollace V, Nottet HS, Clayette P, Turco MC, Muscoli C, Salvemini D, Perno CF (2001). Oxidative stress and neuroAIDS: triggers, modulators and novel antioxidants. *Trends Neurosci* **24**: 411–416.
- Moses AV, Bloom FE, Pauza CD, Nelson JA (1993). HIV infection of human brain capillary endothelial cells occurs via a CD4/galactosylceramide-independent mechanism. *Proc Natl Acad Sci USA* **90**: 10474–10478.
- Nagasawa T, Hirota S, Tachibana K, Takakura N, Nishikawa S, Kitamura Y, Yoshida N, Kikutani H, Kishimoto T (1996). Defects of B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1. *Nature* **382**: 635–638.
- Nath A, Geiger J (1998). Neurobiological aspects of human immunodeficiency virus infection: neurotoxic mechanisms. *Prog Neurobiol* **54**: 19–33.
- Nath A, Hartloper V, Furer M, Fowke KR (1995). Infection of human fetal astrocytes with HIV-1: viral tropism and the role of cell to cell contact in viral transmission. *J Neuropathol Exp Neurol* **54**: 320–330.
- Navia BA, Cho ES, Petito CK, Price RW (1986). The AIDS dementia complex: II. Neuropathology. *Ann Neurol* **19**: 525–535.
- Nishiyori A, Minami M, Ohtani Y, Takami S, Yamamoto J, Kawaguchi N, Kume T, Akaike A, Satoh M (1998). Localization of fractalkine and CX3CR1 mRNAs in rat brain: does fractalkine play a role in signaling from neuron to microglia? *FEBS Lett* **429**: 167–172.
- Peng H, Huang Y, Erichsen D, Zheng J (2002). HIV-1 gp120 and SDF1- α inhibit neural progenitor cell proliferation: links to HIV-1 associated dementia. Paper presented at The 32rd Annual Meeting of Society for Neuroscience, Orlando, Florida, November 2–7.
- Pereira CF, Middel J, Jansen G, Verhoef J, Nottet HS (2001). Enhanced expression of fractalkine in HIV-1 associated dementia. *J Neuroimmunol* **115**: 168–175.
- Perno CF, Crowe SM, Kornbluth RS (1997). A continuing enigma: the role of cells of macrophage lineage in the development of HIV disease. *J Leukoc Biol* **62**: 1–3.
- Persidsky Y (1999). Model systems for studies of leukocyte migration across the blood-brain barrier. *J Neuro Virol* **5**: 579–590.
- Persidsky Y, Ghorpade A, Rasmussen J, Limoges J, Liu XJ, Stins M, Fiala M, Way D, Kim KS, Witte MH, et al (1999). Microglial and astrocyte chemokines regulate monocyte migration through the blood-brain barrier in human immunodeficiency virus-1 encephalitis. *Am J Pathol* **155**: 1599–1611.
- Pulliam L, Herndier BG, Tang NM, McGrath MS (1991). Human immunodeficiency virus-infected macrophages produce soluble factors that cause histological and neurochemical alterations in cultured human brains. *J Clin Invest* **87**: 503–512.
- Ranki A, Nyberg M, Ovod V, Haltia M, Elovaara I, Raininko R, Haapasalo H, Krohn K (1995). Abundant expression of HIV Nef and Rev proteins in brain astrocytes in vivo is associated with dementia. *AIDS* **9**: 1001–1008.
- Ransohoff R (1997). Chemokines in neurological disease models: correlation between chemokine expression

- patterns and inflammatory pathology. *J Leukoc Biol* **62**: 645–652.
- Ransohoff RM (1998). Chemokines and CNS Inflammation. *Neurotransm* **14**: 3–12.
- Rappaport J, Joseph J, Croul S, Alexander G, Del Valle L, Amini S, Khalili K (1999). Molecular pathway involved in HIV-1-induced CNS pathology: role of viral regulatory protein, Tat. *J Leukoc Biol* **65**: 458–465.
- Raychaudhuri SP, Jiang WY, Farber EM (2001). Cellular localization of fractalkine at sites of inflammation: antigen-presenting cells in psoriasis express high levels of fractalkine. *Br J Dermatol* **144**: 1105–1113.
- Rocamora N, Pascual M, Acsady L, Lecea LD, Freund TF (1996). Expression of NGF and NT3 mRNAs hippocampal interneurons innervated by the GABAergic septo-hippocampal pathway. *J Neurosci* **16**: 3991–4004.
- Ruth JH, Volin MV, Haines GK 3rd, Woodruff DC, Katschke KJ, Jr, Woods JM, Park CC, Morel JC, Koch AE (2001). Fractalkine, a novel chemokine in rheumatoid arthritis and in rat adjuvant-induced arthritis. *Arthritis Rheum* **44**: 1568–1581.
- Saad B, Constam D, Ortmann R, Moos M, Fontana A, Schachner M (1991). Astrocyte-derived TGF- β 2 and NGF differentially regulate neural recognition molecule expression by cultured astrocytes. *J Cell Biol* **115**: 473–484.
- Sanders VJ, Pittman CA, White MG, Wang G, Wiley CA, Achim CL (1998). Chemokines and receptors in HIV encephalitis. *AIDS* **12**: 1021–1026.
- Schwaeble WJ, Stover CM, Schall TJ, Dairaghi DJ, Trinder PK, Linington C, Iglesias A, Schubart A, Lynch NJ, Weihe E, Schafer MK (1998). Neuronal expression of fractalkine in the presence and absence of inflammation. *FEBS Lett* **439**: 203–207.
- Sebire G, Emilie D, Wallon C, Hery C, Devergne O, Delfraissy J-F, Galanaud P, Tardieu M (1993). In vitro production of IL-6, IL-1 β , and tumor necrosis factor- α by human embryonic microglial and neural cells. *J Immunol* **150**: 1517–1523.
- Shieh JTC, Albright AV, Sharron M, Gartner S, Strizki J, Doms RW, Gonzalez-Scarano F (1998). Chemokine receptor utilization by human immunodeficiency virus type 1 isolates that replicate in microglia. *J Virol* **72**: 4243–4249.
- Strizki JM, Albright AV, Sheng H, O'Connor M, Perrin L, Gonzalez-Scarano F (1996). Infection of primary human microglia and monocyte-derived macrophages with human immunodeficiency virus type 1 isolates: evidence of differential tropism. *J Virol* **70**: 7654–7662.
- Tong N, Perry SW, Zhang Q, James HJ, Guo H, Brooks A, Bal H, Kinnear SA, Fine S, Epstein LG, et al (2000). Neuronal fractalkine expression in HIV-1 encephalitis: roles for macrophage recruitment and neuroprotection in the central nervous system. *J Immunol* **164**: 1333–1339.
- Tornatore C, Nath A, Amemiya K, Major EO (1991). Persistent human immunodeficiency virus type 1 infection in human fetal glial cells reactivated by T-cell factor(s) or by the cytokines tumor necrosis factor alpha and interleukin-1 beta. *J Virol* **65**: 6094–6100.
- Tsou CL, Haskell CA, Charo IF (2001). Tumor necrosis factor-alpha-converting enzyme mediates the inducible cleavage of fractalkine. *J Biol Chem* **276**: 44622–44626.
- Vallat A-V, Girolami UD, He J, Mhashikar A, Marasco W, Shi B, Gray F, Bell J, Keohane C, Smith TW, Gabuzda D (1998). Localization of HIV-1 co-receptors CCR5 and CXCR4 in the brain of children with AIDS. *Am J Pathol* **152**: 167–178.
- van der Meer P, Ulrich AM, Gonzalez-Scarano F, Lavi E (2000). Immunohistochemical analysis of CCR2, CCR3, CCR5, and CXCR4 in the human brain: potential mechanisms for HIV dementia. *Exp Mol Pathol* **69**: 192–201.
- Vicenzi E, Alfano M, Ghezzi S, Gatti A, Veglia F, Lazzarin A, Sozzani S, Mantovani A, Poli G (2000). Divergent regulation of HIV-1 replication in PBMC of infected individuals by CC chemokines: suppression by RANTES, MIP-1alpha, and MCP-3, and enhancement by MCP-1. *J Leukoc Biol* **68**: 405–412.
- Volin MV, Woods JM, Amin MA, Connors MA, Harlow LA, Koch AE (2001). Fractalkine: a novel angiogenic chemokine in rheumatoid arthritis. *Am J Pathol* **159**: 1521–1530.
- Wiley CA (1995). Quantitative neuropathologic assessment of HIV-1 encephalitis. *Curr Top Microbiol Immunol* **202**: 55–61.
- Wiley CA, Achim C (1994). Human immunodeficiency virus encephalitis is the pathological correlate of dementia in acquired immunodeficiency syndrome. *Ann Neurol* **36**: 673–676.
- Wiley CA, Masliah E, Morey M, Lemere C, Teresa R, Grafe M, Hansen L, Terry R (1991). Neocortical damage during HIV infection. *Ann Neurol* **29**: 651–657.
- Wiley CA, Schrier RD, Nelson JA, Lampert PW, Oldstone MBA (1986). Cellular localization of human immunodeficiency virus infection within the brains of acquired immune deficiency syndrome patients. *Proc Natl Acad Sci USA* **83**: 7089–7093.
- Wu DT, Woodman SE, Weiss JM, McManus CM, D'Aversa TG, Hesselgesser J, Major EO, Nath A, Berman JW (2000). Mechanisms of leukocyte trafficking into the CNS. *J NeuroVirol* **6(Suppl 1)**: S82–S85.
- Yoshida H, Imaizumi T, Fujimoto K, Matsuo N, Kimura K, Cui X, Matsumiya T, Tanji K, Shibata T, Tamo W, et al (2001). Synergistic stimulation, by tumor necrosis factor-alpha and interferon-gamma, of fractalkine expression in human astrocytes. *Neurosci Lett* **303**: 132–136.
- Zheng J, Bauer M, Cotter RL, Ryan LA, Lopez A, Williams C, Ghorpade A, Gendelman HE (2000). Fractalkine mediated macrophage activation by neuronal injury: relevance for HIV-1 associated dementia. Paper presented at The 30th Annual Meeting of Society for Neuroscience, New Orleans, Louisiana, November 4–9.
- Zheng J, Gendelman HE (1997). The HIV-1 associated dementia complex: a metabolic encephalopathy fueled by viral replication in mononuclear phagocytes. *Curr Opin Neurol* **10**: 319–325.
- Zheng J, Lopez A, Erichsen D, Peng H, Bauer M, Williams C, Morgello S, Ghorpade A, Gendelman HE, Cotter RL, Ryan LA (2002). Neuronal fractalkine: regulation of macrophage activation and inflammatory factor production during HIV-1 associated dementia. *J Immunol* Submitted.
- Zheng J, Niemann D, Bauer M, Williams C, Lopez A, Erichsen D, Ryan LA, Cotter RL, Ghorpade A, Swindells S, Gendelman HE (2001). HIV-1 glia interactions in interleukin-8 and growth-related oncogene a secretion, neuronal signaling and demise: relevance for HIV-1-associated dementia. Paper presented at The 8th Conference on Retroviruses and Opportunistic Infections, Foundation for Retrovirology and Human Health, Chicago, Illinois, February 4–8.

- Zheng J, Thylin M, Ghorpade A, Xiong H, Persidsky Y, Cotter R, Niemann D, Che M, Zeng Y, Gelbard H, *et al* (1999). Intracellular CXCR4 signaling, neuronal apoptosis and neuropathogenic mechanisms of HIV-1-associated dementia. *J Neuroimmunol* **98**: 185–200.
- Zlotnik A, Yoshie O (2000). Chemokines: a new classification system and their role in immunity. *Immunity* **12**: 121–127.
- Zou YR, Kottmann AH, Kuroda M, Taniuchi I, Littman DR (1998). Function of the chemokine receptor CXCR4 in haematopoiesis and in cerebellar development. *Nature* **393**: 595–599.
- Zujovic V, Benavides J, Vige X, Carter C, Taupin V (2000). Fractalkine modulates TNF-alpha secretion and neurotoxicity induced by microglial activation [In Process Citation]. *Glia* **29**: 305–315.